

# MA 1101 : Mathematics I

Satvik Saha, 19MS154

October 27, 2019

## Solution 1.

(i) Let  $S \subseteq \mathbb{R}$  be a finite set with  $n \in \mathbb{N}$  elements. We claim that  $S$  has no limit points. We enumerate the elements of  $S$  as  $x_1, x_2, \dots, x_n$ . Let  $a \in \mathbb{R}$ .

(a) If  $a \notin S$ , let us choose  $|x_i - a| > \epsilon_i > 0$ , for all  $i = 1, 2, \dots, n$ . We set  $A_i = (a - \epsilon_i, a + \epsilon_i)$  to be the  $\epsilon_i$  neighbourhood of  $a$ . If  $x_i > a$ , we have  $x_i = a + (x_i - a) > a + \epsilon_i$ , and if  $x_i < a$ , we have  $x_i = a - (a - x_i) < a - \epsilon_i$ . Thus,  $x_i \notin A_i$ .

We set  $A = \bigcap A_i$ . Since  $A$  is the intersection of a finite number of open intervals,  $A$  is also an open interval.

Thus,  $x_i \notin A$  for all  $x_i \in S$ , i.e.  $S \cap A = \emptyset$ . Thus, there is no  $x \in S$  within the  $\epsilon = \min \epsilon_i > 0$  neighbourhood of  $a$ . Hence,  $a$  is not a limit point.

(b) If  $a \in S$ , without loss of generality, we set  $a = x_1$ . We again choose  $|x_i - a| > \epsilon_i > 0$ , for all  $i = 2, 3, \dots, n$ . We set  $A_i = (a - \epsilon_i, a + \epsilon_i)$  to be the  $\epsilon_i$  neighbourhood of  $a$ . Clearly,  $a = x_1 \in A_1$ . Arguing as before,  $x_i \notin A_i$  for  $i = 2, 3, \dots, n$ .

We set  $A = \bigcap A_i$ . Thus,  $a = x_1 \in A$  and  $x_i \notin A$  for  $i \neq 1$ , i.e.  $S \cap A = \{a\}$ . Thus, the only  $x \in S$  within the  $\epsilon = \min \epsilon_i$  neighbourhood of  $a$  is  $a$ . Hence,  $a$  is not a limit point.

Therefore, any finite set  $S$  has no limit points. □

(ii) Let  $S = (0, \infty) \subseteq \mathbb{R}$ . We claim that  $[0, \infty)$  is the set of all limit points of  $S$ . Let  $a \in \mathbb{R}$ .

(a) If  $a \in [0, \infty)$ , let  $\epsilon > 0$  be given. Thus,  $a \geq 0 \Rightarrow a + \epsilon/2 > 0$ , and  $a - \epsilon < a + \epsilon/2 < a + \epsilon$ . Hence, we have found  $x = a + \epsilon/2 \in S$  such that  $x \in (a - \epsilon, a + \epsilon)$  and  $x \neq a$ . Thus,  $a$  is a limit point.

(b) If  $a \notin [0, \infty)$ , i.e.  $a < 0$ , we choose  $\epsilon = -a$ . Hence,  $(a - \epsilon, a + \epsilon) \cap S = (2a, 0) \cap (0, \infty) = \emptyset$ . Thus,  $a$  is not a limit point.

This proves our claim. □

(iii) Let  $S = [1, 2) \cup \{3\}$ . We claim that  $[1, 2]$  is the set of all limit points of  $S$ . Let  $a \in \mathbb{R}$ .

(a) If  $a \in [1, 2)$ , let  $\epsilon > 0$  be given. We set  $\epsilon' = \min\{\epsilon, a - 1, 2 - a\}$ , and  $x = a + \epsilon'/2$ . Thus,  $x > a \geq 1$  and  $x < a + \epsilon' \leq a + (2 - a) = 2$ . Also,  $-\epsilon < \epsilon'/2 < \epsilon$ . Hence, we have found  $x \in (1, 2) \subset S$  such that  $x \in (a - \epsilon, a + \epsilon)$  and  $x \neq a$ . Thus,  $a$  is a limit point.

(b) If  $a \in \{2\}$ , i.e.  $a = 2$ , let  $\epsilon > 0$  be given. We set  $\epsilon' = \min\{\epsilon, 1\}$ , and  $x = a - \epsilon'/2$ . Thus,  $x > a - \epsilon' \geq a - 1 = 1$  and  $x < a = 2$ . Also,  $-\epsilon < -\epsilon'/2 < \epsilon$ . Hence, we have found  $x \in (1, 2) \subset S$  such that  $x \in (a - \epsilon, a + \epsilon)$  and  $x \neq a$ . Thus,  $a$  is a limit point.

(c) If  $a \in \{3\}$ , i.e.  $a = 3$ , we choose  $\epsilon = 1/2 > 0$ . Hence,  $(a - \epsilon, a + \epsilon) \cap S = (2.5, 3.5) \cap ([1, 2) \cup \{3\}) = \{3\}$ . Hence,  $x \in S$  and  $x \in (a - \epsilon, a + \epsilon)$  forces  $x = a$ . Thus,  $a$  is not a limit point.

(d) If  $a < 1$ , we choose  $\epsilon = 1 - a$ . Hence,  $(a - \epsilon, a + \epsilon) \cap S = (2a - 1, 1) \cap ([1, 2) \cup \{3\}) = \emptyset$ . Thus,  $a$  is not a limit point.

(e) If  $2 < a < 3$ , we choose  $\epsilon = \frac{1}{2} \min\{a - 2, 3 - a\}$ . Thus,  $a - \epsilon > a - 2\epsilon \geq a - (a - 2) = 2$  and  $a + \epsilon < a + 2\epsilon \leq a + (3 - a) = 3$ . Therefore,  $(a - \epsilon, a + \epsilon) \subset (2, 3)$ . Hence,  $(a - \epsilon, a + \epsilon) \cap S = \emptyset$ . Thus,  $a$  is not a limit point.

(f) If  $a > 3$ , we choose  $\epsilon = a - 3$ . Hence,  $(a - \epsilon, a + \epsilon) \cap S = (3, 2a - 3) \cap S = \emptyset$ . Thus,  $a$  is not a limit point.

This proves our claim. □

(iv) Let  $S = [1, 2) \cup (2, 3)$ . We claim that  $[1, 3]$  is the set of all limit points of  $S$ . Let  $a \in \mathbb{R}$ .

- (a) If  $a \in (1, 3)$ , let  $\epsilon > 0$  be given. We set  $\epsilon' = \min\{\epsilon, a - 1, 3 - a\}$ , and  $x_- = a - \epsilon'/2$ ,  $x_+ = a + \epsilon'/2$ . Thus,

$$x_- > a - \epsilon' \geq a - (a - 1) = 1,$$

$$x_- < a \leq 3,$$

$$x_+ > a \geq 1,$$

$$x_+ < a + \epsilon' \leq a + (3 - a) = 3.$$

Thus,  $x_-, x_+ \in (1, 3)$ . Since  $x_- < x_+$ , at least one of them is  $x \neq 2$ . Also,  $-\epsilon < -\epsilon'/2 < \epsilon'/2 < \epsilon$ . Hence, we have found  $x \in (1, 3) \setminus \{2\} \subset S$  such that  $x \in (a - \epsilon, a + \epsilon)$  and  $x \neq a$ . Thus,  $a$  is a limit point.

- (b) If  $a \in \{1\}$ , i.e.  $a = 1$ , let  $\epsilon > 0$  be given. We set  $\epsilon' = \min\{\epsilon, 1\}$ , and  $x = a + \epsilon'/2$ . Thus,  $x > a = 1$  and  $x < a + \epsilon' \leq a + 1 = 2$ . Also,  $-\epsilon < \epsilon'/2 < \epsilon$ . Hence, we have found  $x \in (1, 2) \subset S$  such that  $x \in (a - \epsilon, a + \epsilon)$  and  $x \neq a$ . Thus,  $a$  is a limit point.
- (c) If  $a \in \{3\}$ , i.e.  $a = 3$ , let  $\epsilon > 0$  be given. We set  $\epsilon' = \min\{\epsilon, 1\}$ , and  $x = a - \epsilon'/2$ . Thus,  $x > a - \epsilon' \geq a - 1 = 2$  and  $x < a = 3$ . Also,  $-\epsilon < -\epsilon'/2 < \epsilon$ . Hence, we have found  $x \in (2, 3) \subset S$  such that  $x \in (a - \epsilon, a + \epsilon)$  and  $x \neq a$ . Thus,  $a$  is a limit point.
- (d) If  $a < 1$ , we choose  $\epsilon = 1 - a$ . Hence,  $(a - \epsilon, a + \epsilon) \cap S = (2a - 1, 1) \cap S = \emptyset$ . Thus,  $a$  is not a limit point.
- (e) If  $a > 3$ , we choose  $\epsilon = a - 3$ . Hence,  $(a - \epsilon, a + \epsilon) \cap S = (3, 2a - 3) \cap S = \emptyset$ . Thus,  $a$  is not a limit point.

This proves our claim.  $\square$

- (v) Let  $S = \{\frac{1}{n} : n \in \mathbb{N}\}$ . We claim that 0 is the only limit point of  $S$ . Let  $a \in \mathbb{R}$ .

- (a) If  $a = 0$ , let  $\epsilon > 0$  be given. By the *Archimedean Property* of the reals, we choose  $n \in \mathbb{N}$  such that  $n\epsilon > 1$ . Thus,  $\frac{1}{n} \in S$  and  $\frac{1}{n} \in (0 - \epsilon, 0 + \epsilon)$ . Thus, 0 is a limit point.
- (b) If  $a \geq 1$ , we choose  $\epsilon = a - 1$ . Thus,  $(a - \epsilon, a + \epsilon) \cap S = (1, 2a - 1) \cap S = \emptyset$ , since  $S \subset (0, 1]$ . Thus,  $a$  is not a limit point.
- (c) If  $a \in S \setminus \{1\}$ , we find  $n \in \mathbb{N}$  such that  $a = \frac{1}{n}$ . We choose  $\frac{1}{n} - \frac{1}{n+1} > \epsilon > 0$ . Thus,  $a - \epsilon > \frac{1}{n+1}$  and  $a + \epsilon = \frac{2}{n} - \frac{1}{n+1} < \frac{1}{n-1}$ , since  $n^2 - 1 < n^2$ . Hence,  $S \cap (a - \epsilon, a + \epsilon) = \{a\}$ . Thus,  $a$  is not a limit point of  $S$ .
- (d) If  $a \in (0, 1] \setminus S$ , we find  $n \in \mathbb{N}$  such that  $\frac{1}{n+1} < a < \frac{1}{n}$ . We choose  $\min\{\frac{1}{n} - a, a - \frac{1}{n+1}\} > \epsilon > 0$ . Thus,  $a - \epsilon > a - (a - \frac{1}{n+1}) = \frac{1}{n+1}$  and  $a + \epsilon < a + (\frac{1}{n} - a) = \frac{1}{n}$ . Hence,  $S \cap (a - \epsilon, a + \epsilon) = \emptyset$ . Thus,  $a$  is not a limit point.
- (e) If  $a < 0$ , we choose  $\epsilon = -a$ . Hence,  $S \cap (a - \epsilon, a + \epsilon) = S \cap (2a, 0) = \emptyset$ .

Thus proves our claim.  $\square$

- (vi) Let  $S = \{\frac{1}{m} + \frac{1}{n} : m, n \in \mathbb{N}\}$ . We claim that  $\{0\} \cup \{\frac{1}{n} : n \in \mathbb{N}\}$  is the set of all limit points of  $S$ . Let  $a \in \mathbb{R}$ ,  $S' = \{\frac{1}{n} : n \in \mathbb{N}\}$ .

- (a) If  $a = 0$ , let  $\epsilon > 0$  be given. We choose  $n \in \mathbb{N}$  such that  $n\epsilon > 2$ . Thus,  $\frac{2}{n} = \frac{1}{n} + \frac{1}{n} \in S$  and  $\frac{1}{n} + \frac{1}{n} \in (0 - \epsilon, 0 + \epsilon)$ . Thus, 0 is a limit point.
- (b) If  $a \in S'$ , let  $\epsilon > 0$  be given. We find  $n \in \mathbb{N}$  such that  $a = \frac{1}{n}$ . We choose  $k \in \mathbb{N}$  such that  $k\epsilon > 1$ . Thus,  $\frac{1}{n} + \frac{1}{k} \in S$  and  $a < \frac{1}{n} + \frac{1}{k} < \frac{1}{n} + \epsilon$ , so  $\frac{1}{n} + \frac{1}{k} \in (a - \epsilon, a + \epsilon)$ . Thus,  $a$  is a limit point.
- (c) If  $a \notin S'$ ,  $a > 0$ , we choose an  $\epsilon > 0$  such that  $S' \cap (a - \epsilon, a + \epsilon) = \emptyset$ . We can do so since  $a$  is not a limit point of  $S'$ . Also, minimize  $\epsilon$  such that  $a - \epsilon > 0$ .

Consider the elements  $x = \frac{1}{m} + \frac{1}{n} \in S \cap (a - \epsilon/2, a + \epsilon/2)$ , where  $m, n \in \mathbb{N}$ . Without loss of generality, let  $m \leq n$ . Thus,

$$a - \frac{\epsilon}{2} < \frac{1}{n} + \frac{1}{m} < a + \frac{\epsilon}{2}$$

Since  $(a - \epsilon, a + \epsilon)$  has no element of the form  $\frac{1}{k}$  where  $k \in \mathbb{N}$ ,

$$\frac{1}{n} \leq \frac{1}{m} \leq a - \epsilon$$

Also,

$$a - \frac{\epsilon}{2} < \frac{1}{n} + \frac{1}{m} \leq \frac{2}{m}$$

Thus,

$$\frac{1}{m} > \frac{1}{2}(a - \frac{\epsilon}{2})$$

This means that there are only a finite number of  $m$ . Also,

$$a - \frac{\epsilon}{2} < \frac{1}{n} + \frac{1}{m} < \frac{1}{n} + a - \epsilon$$
$$\frac{1}{n} > \frac{\epsilon}{2}$$

Thus, there are only a finite number of  $n$ . This means that there are a finite number of  $x$ .

Hence,  $S \cap (a - \epsilon/2, a + \epsilon/2)$  is a finite set. Hence,  $a$  is not a limit point.

(d) If  $a < 0$ , we choose  $\epsilon = -a$ . Hence,  $S \cap (a - \epsilon, a + \epsilon) = S \cap (2a, 0) = \emptyset$ .

This proves our claim. □

**Solution 2.** Note that for any  $x \in \mathbb{R}$ ,  $x$  is trivially a limit point of  $\mathbb{R}$ , since every  $\epsilon > 0$  neighbourhood of  $\mathbb{R}$  contains infinitely many real numbers other than  $x$ . In addition, removing a finite number of points from  $\mathbb{R}$  means that  $x$  is still a limit point of  $\mathbb{R}$ .

(i) We have  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) := \lfloor x \rfloor$ . We claim that  $\lim_{x \rightarrow 0} f(x)$  does not exist.

Suppose not, i.e.  $\lim_{x \rightarrow 0} f(x) = L$ . We find  $\delta$  such that

$$0 < |x - 0| < \delta \implies |f(x) - L| < \frac{1}{4}$$

We choose  $0 < x_0 < \min\{1, \delta\}$ . Thus,  $f(x_0) - f(-x_0) = 1$ . Now,

$$\begin{aligned} 1 &= |f(x_0) - f(-x_0)| \\ &= |(f(x_0) - L) - (f(-x_0) - L)| \\ &\leq |f(x_0) - L| + |f(-x_0) - L| \\ &< \frac{1}{4} + \frac{1}{4} \\ &= \frac{1}{2} \end{aligned}$$

This is a contradiction, thus proving our claim. □

(ii) We have  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) := \lfloor x \rfloor - \lfloor x/3 \rfloor$ . We claim that  $\lim_{x \rightarrow 0} f(x) = 0$ .

Let  $\epsilon > 0$  be given. We set  $\delta = \frac{1}{2}$ .

Then, for all  $x \in \mathbb{R}$  satisfying  $0 < |x - 0| < \delta$ , we have  $|\lfloor x \rfloor - \lfloor x/3 \rfloor - 0| = 0 < \epsilon$ . This proves our claim. □

(iii) We have  $f: \mathbb{R} \setminus \{2\} \rightarrow \mathbb{R}$ ,  $f(x) = \frac{x^3 - 8}{x - 2}$ . We claim that  $\lim_{x \rightarrow 2} f(x) = 12$ .

Let  $\epsilon > 0$  be given. We set  $\delta = \min\{1, \epsilon/7\}$ .

Then, for all  $x \in \mathbb{R} \setminus \{2\}$  satisfying  $0 < |x - 2| < \delta$ , we have

$$\begin{aligned} \left| \frac{x^3 - 8}{x - 2} - 12 \right| &= |x^2 + 2x + 4 - 12| \\ &= |x^2 + 2x - 8| \\ &= |(x - 2)(x + 4)| \\ &= |x - 2| |x - 2 + 6| \\ &\leq |x - 2| (|x - 2| + 6) \\ &< \delta(\delta + 6) \\ &\leq \frac{\epsilon}{7}(1 + 6) \\ &= \epsilon \end{aligned}$$

This proves our claim. □

(iv) We have  $f: \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ ,  $f(x) := x \sin \frac{1}{x}$ . We claim that  $\lim_{x \rightarrow 0} f(x) = 0$ .

Let  $\epsilon > 0$  be given. We set  $\delta = \epsilon$ .

Then, for all  $x \in \mathbb{R} \setminus \{0\}$  satisfying  $0 < |x - 0| < \delta$ , we have  $|x \sin \frac{1}{x}| \leq |x| < \epsilon$ . This proves our claim.  $\square$

(v) We have  $f: \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ ,  $f(x) := x/|x|$ . We claim that  $\lim_{x \rightarrow 0} f(x)$  does not exist.

Suppose not, i.e.  $\lim_{x \rightarrow 0} f(x) = L$ . We find  $\delta$  such that

$$0 < |x - 0| < \delta \implies |f(x) - L| < \frac{1}{2}$$

Note that  $f(x) - f(-x) = 2$ . Thus,

$$\begin{aligned} 2 &= |f(\delta/2) - f(-\delta/2)| \\ &= |(f(\delta/2) - L) - (f(-\delta/2) - L)| \\ &\leq |f(\delta/2) - L| + |f(-\delta/2) - L| \\ &< \frac{1}{2} + \frac{1}{2} \\ &= 1 \end{aligned}$$

This is a contradiction, thus proving our claim.  $\square$

**Solution 3.** Let  $\emptyset \neq D \subseteq \mathbb{R}$ ,  $f, g: D \rightarrow \mathbb{R}$  and let  $a$  be a limit point of  $D$ . Let  $\lim_{x \rightarrow a} f(x)$  and  $\lim_{x \rightarrow a} g(x)$  exist. We write

$$\lim_{x \rightarrow a} f(x) := L, \quad \lim_{x \rightarrow a} g(x) := M.$$

(i) We claim that  $\lim_{x \rightarrow a} (f(x) + g(x)) = L + M$ .

Let  $\epsilon > 0$  be given. We find  $\delta_f, \delta_g$  such that for all  $x \in D$ ,

$$0 < |x - a| < \delta_f \implies |f(x) - L| < \epsilon/2,$$

$$0 < |x - a| < \delta_g \implies |g(x) - M| < \epsilon/2.$$

We set  $\delta = \min\{\delta_f, \delta_g\}$ . Then, for all  $x \in D$  satisfying  $0 < |x - a| < \delta$ , we have

$$\begin{aligned} |(f(x) + g(x)) - (L + M)| &= |(f(x) - L) + (g(x) - M)| \\ &\leq |f(x) - L| + |g(x) - M| \\ &< \epsilon/2 + \epsilon/2 \\ &= \epsilon \end{aligned}$$

This proves our claim.  $\square$

(ii) We claim that for all  $\alpha \in \mathbb{R}$ ,  $\lim_{x \rightarrow a} (\alpha f(x)) = \alpha L$ .

Let  $\epsilon > 0$  be given. If  $\alpha \neq 0$ , we find  $\delta_f$  such that for all  $x \in D$ ,

$$0 < |x - a| < \delta_f \implies |f(x) - L| < \epsilon/|\alpha|.$$

We set  $\delta = \delta_f$ . Then, for all  $x \in D$  satisfying  $0 < |x - a| < \delta$ , we have

$$\begin{aligned} |\alpha f(x) - \alpha L| &= |\alpha| |f(x) - L| \\ &< |\alpha| \frac{\epsilon}{|\alpha|} \\ &= \epsilon \end{aligned}$$

If  $\alpha = 0$ , we trivially have

$$0 < |x - a| < \delta = \epsilon \implies |\alpha f(x) - \alpha L| = 0 < \epsilon.$$

This proves our claim.  $\square$

(iii) We claim that  $\lim_{x \rightarrow a} f(x)g(x) = LM$ . To prove this, we first show that  $\lim_{x \rightarrow a} (f(x) - L)(g(x) - M) = 0$ .

Let  $\epsilon > 0$  be given. We find  $\delta_f, \delta_g$  such that for all  $x \in D$ ,

$$0 < |x - a| < \delta_f \implies |f(x) - L| < \sqrt{\epsilon},$$

$$0 < |x - a| < \delta_g \implies |g(x) - M| < \sqrt{\epsilon}.$$

We set  $\delta = \min\{\delta_f, \delta_g\}$ . Then, for all  $x \in D$  satisfying  $0 < |x - a| < \delta$ , we have

$$\begin{aligned} |(f(x) - L)(g(x) - M) - 0| &= |f(x) - L||g(x) - M| \\ &< \sqrt{\epsilon}\sqrt{\epsilon} \\ &= \epsilon \end{aligned}$$

Thus,  $\lim_{x \rightarrow a} (f(x) - L)(g(x) - M) = 0$ .

We now show that for a constant function  $h: D \rightarrow \mathbb{R}$ ,  $h(x) = k$ , we have  $\lim_{x \rightarrow a} h(x) = k$ .

Let  $\epsilon > 0$  be given. We set  $\delta = \epsilon$ . Then, for all  $x \in D$  satisfying  $0 < |x - a| < \delta$ , we have  $|h(x) - k| = 0 < \epsilon$ .

Therefore,

$$\begin{aligned} 0 &= \lim_{x \rightarrow a} (f(x) - L)(g(x) - M) \\ &= \lim_{x \rightarrow a} (f(x)g(x) - Lg(x) - Mf(x) + LM) \\ &= \lim_{x \rightarrow a} f(x)g(x) - \lim_{x \rightarrow a} Lg(x) - \lim_{x \rightarrow a} Mf(x) + \lim_{x \rightarrow a} LM \\ &= \lim_{x \rightarrow a} f(x)g(x) - L \lim_{x \rightarrow a} g(x) - M \lim_{x \rightarrow a} f(x) + LM \\ &= \lim_{x \rightarrow a} f(x)g(x) - LM - ML + LM \\ &= \lim_{x \rightarrow a} f(x)g(x) - LM \end{aligned}$$

$$\lim_{x \rightarrow a} f(x)g(x) = LM$$

□

(iv) We claim that if  $M \neq 0$ ,  $\lim_{x \rightarrow a} f(x)/g(x) = L/M$ . To prove this, we first show that  $\lim_{x \rightarrow a} 1/g(x) = 1/M$ .

Let  $\epsilon > 0$  be given. We find  $\delta_1, \delta_2$  such that for all  $x \in D$ ,

$$0 < |x - a| < \delta_1 \implies |g(x) - M| < \frac{1}{2}|M|,$$

$$0 < |x - a| < \delta_2 \implies |g(x) - M| < \frac{1}{2}\epsilon|M|^2.$$

We set  $\delta = \min\{\delta_1, \delta_2\}$ . Then, for all  $x \in D$  satisfying  $0 < |x - a| < \delta$ , we have

$$\begin{aligned}\frac{1}{2}|M| &> |g(x) - M| \\ &\geq ||g(x)| - |M|| \\ &\geq |M| - |g(x)| \\ |g(x)| &> \frac{1}{2}|M| > 0 \\ \frac{1}{|g(x)|} &< \frac{2}{|M|} \\ \left| \frac{1}{g(x)} - \frac{1}{M} \right| &= \frac{|g(x) - M|}{|Mg(x)|} \\ &= |g(x) - M| \frac{1}{|M||g(x)|} \\ &< \frac{1}{2}\epsilon|M|^2 \frac{2}{|M|^2} \\ &= \epsilon\end{aligned}$$

Thus,  $\lim_{x \rightarrow a} 1/g(x) = 1/M$ . Therefore,

$$\begin{aligned}\lim_{x \rightarrow a} f(x)g(x) &= \lim_{x \rightarrow a} f(x) \frac{1}{g(x)} \\ &= \lim_{x \rightarrow a} f(x) \lim_{x \rightarrow a} \frac{1}{g(x)} \\ &= \frac{L}{M}\end{aligned}$$

□